

Clinical Policy: Esketamine (Spravato)

Reference Number: NH.PMN.199

Effective Date: 06.24

Last Review Date: 06.24

Line of Business: Medicaid

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Esketamine (Spravato™) is a non-competitive N-methyl D-aspartate (NMDA) receptor antagonist.

FDA Approved Indication(s)

Spravato, in conjunction with an oral antidepressant, is indicated for the treatment of:

- Treatment-resistant depression (TRD) in adults.
- Depressive symptoms in adults with major depressive disorder (MDD) with acute suicidal ideation or behavior.

Limitation(s) of use:

- Spravato is not approved as an anesthetic agent. The safety and effectiveness of Spravato as an anesthetic agent have not been established.
- The effectiveness of Spravato in preventing suicide or in reducing suicidal ideation or behavior has not been demonstrated. Use of Spravato does not preclude the need for hospitalization if clinically warranted, even if patients experience improvement after an initial dose of Spravato.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation® that Spravato is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Treatment-Resistant Depression (must meet all):

1. Diagnosis of TRD;
2. Prescribed by or in consultation with a psychiatrist;
3. Age ≥ 18 and < 65 years (*see Appendix E*);
4. Member has a documented baseline Patient Health Questionnaire (PHQ-9) score ≥ 15 , indicating moderately severe major depression, within the previous four weeks (*see Appendix D*);
5. Trial and failure of at least two (2) preferred products unless contraindicated or clinically significant adverse effects are experienced;
6. Currently stabilized on an oral antidepressant for at least two weeks and (a):
 - a. Spravato will be used in combination with oral antidepressant;

CLINICAL POLICY

Esketamine

7. Member meets one of the following (a or b):
 - a. No prior history of treatment with Spravato;
 - b. Documentation of a prior positive response to Spravato as documented by a history of $\geq 50\%$ reduction in PHQ-9 score;
8. Dose does not exceed 168 mg per week during four week induction phase.

Approval duration: 3 months (up to 23 nasal spray devices)

B. Major Depressive Disorder with Suicidal Ideation or Behavior (must meet all):

1. Diagnosis of MDD;
2. Prescribed by or in consultation with a psychiatrist;
3. Age ≥ 18 years;
4. Spravato is prescribed in combination with initiation or optimization of oral antidepressant therapy;
5. Trial and failure of at least two (2) preferred products unless contraindicated or clinically significant adverse effects are experienced;
6. Member is recently (within the last 5 days) discharged from or currently in an acute or subacute inpatient care for suicidality;
7. Member meets one of the following (a or b):
 - a. No prior history of treatment with Spravato;
 - b. Documentation of a prior positive response to Spravato (*see Appendix E*);
8. Member meets one of the following (a, b, or c):
 - a. Montgomery-Åsberg Depression Rating Scale (MADRS) score is ≥ 20 (moderate depression) (*see Appendix D*);
 - b. Hamilton Rating Scale for Depression (HAM-D) score is ≥ 17 (moderate depression) (*see Appendix D*);
 - c. PHQ-9 score is ≥ 15 (moderately severe depression) (*see Appendix D*);
9. Dose does not exceed 84 mg (3 nasal spray devices) twice weekly.

Approval duration: 4 weeks (up to 23 nasal spray devices)

C. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the PDL, the no coverage criteria policy: CP.PMN.255; or
 - b. For drugs NOT on the PDL, the non-formulary policy: CP.PMN.16; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy: CP.PMN.53.

II. Continued Therapy

A. Treatment-Resistant Depression (must meet all):

1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;

CLINICAL POLICY

Esketamine

- b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
2. Member is responding positively to therapy as evidenced by at least a 50% reduction in PHQ-9 score compared to baseline (*see Appendix D*);
3. Spravato is being used in combination with an oral antidepressant;
4. If request is for a dose increase, new dose does not exceed 84 mg (3 nasal spray devices) per week.

Approval duration: 6 months

B. Major Depressive Disorder with Suicidal Ideation or Behavior

1. Re-authorization is not permitted. Members must meet the initial approval criteria.

Approval duration: Not applicable

C. Other diagnoses/indications (must meet 1 or 2):

1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the PDL, the no coverage criteria policy: CP.PMN.255; or
 - b. For drugs NOT on the PDL, the non-formulary policy: CP.PMN.16; or
2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy: CP.PMN.53.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.PMN.53 or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration

HAM-D: Hamilton Rating Scale for Depression

MADRS: Montgomery-Åsberg Depression Rating Scale

MDD: major depressive disorder

NMDA: non-competitive N-methyl D aspartate

PHQ-9: Patient Health Questionnaire

SNRI: serotonin norepinephrine reuptake inhibitor

SSRI: selective serotonin reuptake inhibitor

TCA: tricyclic antidepressant

TRD: treatment-resistant depression

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

CLINICAL POLICY
Esketamine

| Drug Name | Dosing Regimen | Dose Limit/ Maximum Dose |
|---|---|--|
| <i>SSRI</i> | | |
| citalopram (Celexa [®]) | 20 mg PO QD; may increase to 40 mg PO QD after one week | 40 mg/day (≤ 60 years) 20 mg/day (> 60 years) |
| escitalopram (Lexapro [®]) | 10 mg PO QD; may increase to 20 mg PO QD after 1 week | 20 mg/day |
| fluoxetine (Prozac [®] , Prozac Weekly [®]) | Prozac: 20 mg PO QD; may increase by 10-20 mg after several weeks Prozac Weekly: 90 mg PO q week beginning 7 days after the last daily dose | Prozac: 80 mg/day Prozac Weekly: 90 mg/week |
| paroxetine (Paxil [®] , Paxil CR [®] , Pexeva [®]) | Paxil, Pexeva: 20 mg PO QD; may increase by 10 mg every week as needed Paxil CR: 25 mg PO QD; may increase by 12.5 mg every week as needed | Paxil, Pexeva: 50 mg/day Paxil CR: 62.5 mg/day |
| sertraline (Zoloft [®]) | 50 mg PO QD; may increase every week as needed | 200 mg/day |
| <i>SNRIs</i> | | |
| duloxetine (Cymbalta [®]) | 20 mg PO BID or 30 mg PO BID or 60 mg PO QD | 120 mg/day |
| venlafaxine (Effexor [®] , Effexor XR [®]) | Effexor: 75 mg/day PO in 2-3 divided doses; may increase by 75 mg every 4 days as needed Effexor XR: 75 mg PO QD; may increase by 75 mg every 4 days as needed | Effexor: 225 mg/day (outpatient) or 375 mg/day (inpatient) Effexor XR: 225 mg/day |
| desvenlafaxine (Pristiq [®] , Khedezla [®]) | 50 mg PO QD | 400 mg/day |
| Fetzima [®] (levomilnacipran) | 20 mg PO QD for 2 days, then 40 mg PO QD; may increase by 40 mg every 2 days | 120 mg/day |
| <i>TCA</i> | | |
| amitriptyline (Elavil [®]) | 50 to 100 mg/day PO QD or divided doses | 150 mg/day |
| amoxapine | 25 to 300 mg/day PO in divided doses | 400 mg/day (300 mg/day if geriatric) |
| clomipramine* (Anafranil [®]) | 12.5 to 150 mg/day PO QD | 250 mg/day (200 mg/day if pediatric) |
| desipramine (Norpramin [®]) | 25 to 300 mg/day PO QD | 300 mg/day (100 mg/day if pediatric) |
| doxepin (Sinequan [®]) | 25 to 300 mg/day PO QD | 300 mg/day |
| imipramine HCl (Tofranil [®]) | 25 to 200 mg/day PO QD or divided doses | 200 mg/day (150 mg/day if geriatric or pediatric) |

CLINICAL POLICY

Esketamine

| Drug Name | Dosing Regimen | Dose Limit/ Maximum Dose |
|--|--|--|
| imipramine pamoate (Tofranil PM [®]) | 25 to 200 mg/day PO QD or divided doses | 200 mg/day (100 mg/day if geriatric or pediatric) |
| nortriptyline (Pamelor [®]) | 25 to 150 mg/day PO QD | 150 mg/day |
| protriptyline (Vivactil [®]) | 10 to 60 mg/day PO in divided doses | 60 mg/day (30 mg/day if geriatric or pediatric) |
| trimipramine (Surmontil [®]) | 25 to 200 mg/day PO QD | 200 mg/day (100 mg/day if geriatric or pediatric) |
| <i>Second Generation Antipsychotics</i> | | |
| aripiprazole (Abilify [®]) | 2 to 15 mg PO QD | 15 mg/day |
| Rexulti [®] (brexpiprazole) | 0.5 to 3 mg PO QD | 3 mg/day |
| Vraylar [®] (cariprazine)* | 0.5 to 4.5 mg PO QD | 4.5 mg/day |
| olanzapine (Zyprexa [®])* | 5 to 20 mg PO QD | 20 mg/day |
| quetiapine (Seroquel [®])* | 25 to 400 mg PO QD | 400 mg/day |
| risperidone (Risperdal [®])* | 0.25 to 3 mg PO QD | 3 mg/day |
| ziprasidone (Geodon [®])* | 20 to 80 mg PO BID | 160 mg/day |
| <i>Other Antidepressants</i> | | |
| bupropion (Aplenzin [®] , Budeprion SR [®] , Budeprion XL [®] , Forfivo XL [®] , Wellbutrin [®] , Wellbutrin SR [®] , Wellbutrin XL [®]) | Varies | Immediate-release: 450 mg/day (300 mg/day if pediatric) Sustained-release: 400 mg/day Extended-release (HCl): 450 mg/day Extended-release (HBr): 522 mg/day |
| bupirone* | 15 to 20 mg/day PO in 2 divided doses | 60 mg/day |
| mirtazapine (Remeron [®]) | 15 to 45 mg PO QD | 45 mg/day |
| lithium* | 300 mg PO QD or BID; up to 600 to 1,200 mg PO daily in divided doses | 1,200 mg/day |
| thyroid hormone* | 25 to 50 mcg/day PO | 50 mcg/day |

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

*Off-label

CLINICAL POLICY

Esketamine

Appendix C: Contraindications/Boxed Warnings

- Spravato is not indicated for the treatment of bipolar depression.
- Contraindication(s):
 - Aneurysmal vascular disease (including thoracic and abdominal aorta, intracranial, and peripheral arterial vessels) or arteriovenous malformation.
 - History of intracerebral hemorrhage.
 - Hypersensitivity to esketamine, ketamine, or any of the excipients.
- Boxed warning(s):
 - Risk for sedation and dissociation after administration. Monitor patients for at least two hours after administration.
 - Potential for abuse and misuse. Consider the risks and benefits of prescribing Spravato prior to using in patients at higher risk of abuse. Monitor patients for signs and symptoms of abuse and misuse.
 - Spravato is only available through a restricted program called the Spravato REMS.
 - Increased risk of suicidal thoughts and behaviors in pediatric and young adult patients taking antidepressants. Closely monitor all antidepressant-treated patients for clinical worsening and emergence of suicidal thoughts and behaviors. Spravato is not approved for use in pediatric patients. Spravato is available only through a restricted program under a REMS called the Spravato REMS because of the risks of serious adverse outcomes from sedation, dissociation, and abuse and misuse.
- Healthcare settings must be certified in the REMS program and ensure that Spravato is:
 - Only dispensed in healthcare settings and administered to patients who are enrolled in the program.
 - Administered by patients under the direct observation of a healthcare provider and that patients are monitored by a healthcare provider for at least 2 hours after administration of Spravato.
 - Pharmacies must be certified in the REMS and must only dispense Spravato to healthcare settings that are certified in the program.
 - Further information, including a list of certified pharmacies is available at www.Spravatorems.com or 1-855-382-6022.

Appendix D: PHQ-9, MADRS, and HAM-D Rating Scales

- The PHQ-9 is a 9-item multiple choice questionnaire used for diagnosis, screening, monitoring and measuring the severity of depression.

| PHQ-9 Score | Depression Severity |
|-------------|--|
| 5 – 9 | Minimal symptoms |
| 10 – 14 | Minor depression Major depression, mild |
| 15 – 19 | Major depression, moderately severe |
| > 20 | Major depression, severe |

- The MADRS is a 10-item diagnostic questionnaire used to measure the severity of depressive episodes in patients with mood disorders.

CLINICAL POLICY

Esketamine

| MADRS Score | Depression Rating |
|-------------|-----------------------|
| 0 – 6 | Normal/symptom absent |
| 7 – 19 | Mild depression |
| 20 – 34 | Moderate depression |
| > 34 | Severe depression |

- The HAM-D17 scale is a 17-item depression assessment scale to assess severity of, and change in, depressive symptoms.

| HAM-D Score | Depression Rating |
|-------------|--|
| 0 – 7 | Normal, absence or remission of depression |
| 8 – 16 | Mild depression |
| 17 – 23 | Moderate depression |
| > 24 | Severe depression |

Appendix E: General Information

- Positive responses to therapy include but are not limited to:
 - Previous demonstrated improvement in depressive symptoms
 - Rapid reduction in depressive symptoms and thus rapid reduction in suicidality, either during hospitalization, or during a previous episode of suicidality.
 - Improvement from baseline in PHQ-9, MADRS, or HAM-D17 score.
- The efficacy of Spravato for the treatment of TRD in geriatric patients was evaluated in a 4-week, randomized, double-blind study with patients receiving placebo or Spravato intranasally plus an oral antidepressant (TRANSFORM-3).
 - The trial included patients between the ages of 65 and 74 years old.
 - At the end of four weeks, Spravato plus antidepressant did not achieve statistically significant difference when compared to those receiving placebo plus antidepressant on the primary efficacy endpoint of change from baseline to Week 4 on the MADRS.
 - During the double-blind phase, TEAEs occurred in 70.8% (51/72) of patients receiving antidepressant plus Spravato and 60.0% (39/65) receiving antidepressant plus placebo. Overall, safety results were consistent with those reported in previous esketamine studies in younger adults, including those in patients \geq 75 years old.

Appendix F: States with Limitations against Redirections in Certain Mental Health Settings

| State | Step Therapy Prohibited? | Notes |
|-------|--------------------------|--|
| TX | No | <i>*Applies to HIM requests only*</i> Failure of ONE of the following antidepressant augmentation therapies, used for \geq 4 weeks, unless clinically significant adverse effects are experienced or all are contraindicated: second-generation antipsychotic, lithium, thyroid hormone, bupirone |

V. Dosage and Administration

| Indication | Dosing Regimen | Maximum Dose |
|------------|--|--------------|
| TRD | Administer in conjunction with an oral antidepressant. | 84 mg/dose |

CLINICAL POLICY

Esketamine

| Indication | Dosing Regimen | Maximum Dose |
|---|--|--------------|
| | Induction Phase <u>Weeks 1 to 4:</u> Administer nasally twice per week Day 1 starting dose: 56 mg Subsequent doses: 56 mg or 84 mg Maintenance Phase <u>Weeks 5 to 8:</u> Administer 56 mg or 84 mg nasally once weekly <u>Week 9 and after:</u> Administer 56 mg or 84 mg every 2 weeks or once weekly | |
| Depressive symptoms with MDD with acute suicidal ideation or behavior | Administer in conjunction with an oral antidepressant. Administer 56 mg or 84 mg nasally twice weekly for 4 weeks. | 84 mg/dose |

VI. Product Availability

Nasal spray: 28 mg of esketamine per device. Each nasal spray device delivers two sprays containing a total of 28 mg esketamine.

VII. References

1. Spravato Prescribing Information. Titusville, NJ: Janssen Pharmaceuticals; July 2020. Available at: <http://www.janssenlabels.com/package-insert/product-monograph/prescribing-information/SPRAVATO-pi.pdf>. Accessed February 7, 2023.
2. American Psychiatric Association. Practice guideline for the treatment of patients with major depressive disorder, third edition. November 2010. Available at: <http://psychiatryonline.org/guidelines.aspx>.
3. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc. Updated periodically. Accessed February 7, 2023.
4. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med.* 2001;16(9):606–613.
5. Montgomery–Åsberg Depression Rating Scale. Available at: http://www.liquisearch.com/montgomery%E2%80%93%C3%85sberg_depression_rating_scale/interpretation. Accessed February 9, 2022.
6. Sharp, Rachel. The Hamilton rating scale for depression. *Occupational Medicine.* 2015; 65(4):340.
7. Ochs-Ross R, Daly EJ, Zhang Y et al. Efficacy and safety of esketamine nasal spray plus an oral antidepressant in elderly patients with treatment-resistant depression TRANSFORM-3. *Am J Geriatr Psychiatry.* 2020 Feb;28(2):121-141.

Coding Implications

CLINICAL POLICY

Esketamine

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

| HCPCS Codes | Description |
|-------------|--|
| G2082 | Office or other outpatient visit for the evaluation and management of an established patient that requires the supervision of a physician or other qualified health care professional and provision of up to 56 mg of esketamine nasal self-administration, includes 2 hours post administration observation |
| G2083 | Office or other outpatient visit for the evaluation and management of an established patient that requires the supervision of a physician or other qualified health care professional and provision of greater than 56 mg esketamine nasal self-administration, includes 2 hours post administration observation |

| Reviews, Revisions, and Approvals | Date | P&T Approval Date |
|-----------------------------------|-------|-------------------|
| Policy created | 06.24 | 06.24 |

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a

CLINICAL POLICY

Esketamine

discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2019 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.